

Healthy Kids Survey  
Sunset High School  
Selected Findings 2017  
N=90<sup>\*†‡</sup>

Date of Survey: Spring 2017

Notes:

---

\* 95% of students in attendance on the survey date


† X% of students enrolled

‡ 11% of collected surveys deemed unreliable




## Gender Distribution

Sunset 2017




## Grade Level Distribution

Sunset 2017




# Ethnicity Distribution

Sunset 2017




# Lifetime Substance Use

Sunset 2017




# Lifetime Smoking

Sunset 2017





# Past Month Substance Use

Sunset 2017



# Past Month Smoking



Sunset 2017



# Number of Binge Drinking Episodes (>4 Drinks/Occasion)


(Past 30 Days)

Sunset 2017



# Ever Been Drunk or High on Drugs on School Property

Sunset 2017



Sunset High School  
Healthy Kids Survey  
2017

**Lifetime Substance Use**

	9th & 10th	11th	12th	Overall
	(%)	(%)	(%)	Mean
	(%)	(%)	(%)	(%)
Alcohol	75.0	76.0	85.7	80.5
Marijuana	75.0	68.0	85.7	77.9
Cigarettes	50.0	53.8	54.3	53.8
Smokeless tobacco	18.8	30.8	22.9	24.4
Hookah	50.0	50.0	57.1	52.6
Vaping devices	56.2	53.8	80.0	66.7
Methamphetamine	31.2	23.1	11.4	19.2
Inhalants	43.8	23.1	14.3	24.4
LSD	43.8	24.0	28.6	31.2
Ecstasy	31.2	12.0	14.3	16.9
Heroin	0.0	0.0	0.0	0.0
Cocaine/crack	31.2	23.1	22.9	25.6
Prescription pain killers	37.5	26.9	28.6	30.8
Barbiturates	18.8	3.8	0.0	5.1
Tranquilizers	37.5	38.5	31.4	34.6
Cough medicines	50.0	30.8	34.3	35.9
Diet pills	18.8	7.7	5.7	9.0
Ritalin or adderall	43.8	23.1	31.4	32.1
Other drugs*	64.7	65.4	57.1	62.0

### Past 30 Days Substance Use

	9th & 10th	11th	12th	Overall
	(%)	(%)	(%)	Mean
	(%)	(%)	(%)	(%)
Alcohol	46.7	40.0	57.1	50.0
Marijuana	53.3	56.0	58.8	57.3
Cigarettes	40.0	36.0	23.5	32.0
Smokeless tobacco	13.3	0.0	2.9	4.1
Hookah	26.7	20.8	17.6	20.3
Vaping devices	46.7	43.5	51.4	48.6
Methamphetamine	0.0	4.3	0.0	1.4
Inhalants	6.7	4.2	5.9	6.8
LSD	14.3	13.0	5.9	11.1
Ecstasy	6.7	0.0	0.0	1.4
Cocaine	7.1	13.0	11.8	11.1
Prescription pain killers	40.0	17.4	14.7	21.9
Two or more drugs together	46.7	26.1	26.5	31.5
Other drugs*	35.3	30.8	20.0	27.5

### Recent Binge Drinking (Past 30 Days)

	9th & 10th	11th	12th	Overall
	(%)	(%)	(%)	Mean
	(%)	(%)	(%)	(%)
All students	46.7	33.3	31.4	36.0
Recent drinkers	100.0	88.9	55.0	73.0

This report was generated on June 9, 2017



## Notes

Students' responses were deemed unreliable if recent substance use history was inconsistent with their lifetime responses or they reported using a fake drug.

Ethnicity Distribution: other signifies mixed (two or more) races

Lifetime Substance Use: other drugs include methamphetamine or any amphetamine, inhalants, LSD or other psychedelics, ecstasy, heroin, cocaine, cough/cold medicines, Ritalin™, or Adderall™, barbiturates, tranquilizers or sedatives, and any other drug or pill to get "high"

Past Month Substance Use: other drugs includes methamphetamine or any amphetamine, inhalants, LSD or other psychedelics, ecstasy, cocaine, prescription pain medication to get "high" or for reasons other than prescribed (such as Vicodin™, OxyContin™, Percodan™, Ritalin™, Adderall™, Xanax™) and any other drug or pill to get "high".

When multiple drugs are tallied in computations, such as when other drugs are reported, the tables report data for the complete cases and the graphs provide reports for all cases. Complete cases are reports where an individual provided a valid response to all of the terms in a related set. Estimates were insensitive to this analytical decision, resulting in variance of 0.5 percentage points or less.